

A NEW VARIETY OF *COELACHNE* R. BR. (POACEAE) FROM INDIA—
COELACHNE PERPUSILLA (ARN. EX STEUD.) THW. VAR.
NILAGIRICA VED PRAKASH ET JAIN VAR. NOV.

Varietati *perpusillae* affinis, sed differt culmis latoribus, vaginis hirsutis, laminis latoribus, spiculis et ramis paniculae majoribus, et lemma paleaque superiore enervi ecarinataque.

Holotypus lectus a J. L. Ellis 34627 ad locum Parthimund, Tamil Nadu, Nilgiri, alt. 2300 m, die July 11, 1970 et Positus in CAL. Isotypus positus in MH.

Perennial, densely tufted ca 15-30 cm tall, ascending, with numerous slender fibrous roots. Culms slender, terete; internodes up to 3.5 cm long, faintly striate, glabrous; nodes hirsute. Leaf-sheaths mostly shorter than the internodes, terete, hirsute; ligules 0 (absent) or minute fringe of hairs; blades 1.0-3.5 cm long, 1.5-4.0 mm broad, linear-lanceolate to lanceolate, acute, subrounded at base, flat, striate and glabrous or scaberrulous above, hirsute beneath, nerves not or slightly distinct beneath.

Inflorescence a panicle 3.5-8.0 cm long, lax; axis subterete to channelled, slightly flexuous, glabrous; branches up to 3 cm long, distant, spreading, slender, sub-terete, glabrous; pedicels (♂) 0.5-7.0 mm long, similar to branches. Spikelets ca 3.5-4.0 mm long, oblong to obovoid-oblong or ovate-oblong, gaping, 2-flowered, somewhat laterally compressed. Glumes 2, ovate-elliptic, obtuse-apiculate, membranous (-hyaline), glabrous; lower glume shorter, 1.7-2.0 mm long, flatter, margins narrowly incurved, 1-3-nerved (lateral nerves only near the base; upper glume longer and broader, 2.3-2.5 mm long, rounded on back, margins widely incurved, 7-nerved. Florets 2, exserted. Lower floret ♂ subsessile, ± 3 mm long, lanceolate, obtuse, glabrous; callus densely bearded, hairs up to 1 mm long; lemma ovate-elliptic, obtuse, folded, coriaceous,

nerveless (-1-nerved), glabrous; palea elliptic obtuse, margins widely incurved, grooved on back, nerveless, glabrous; stamens 3; anthers ± 1.5-1, 7 mm long, oblong; filaments ± 0.2-0.5 mm long; ovary ± 0.5 mm long, ovoid; styles 2, 0.2-0.4 mm long, distinct; stigmas ± 1 mm long, distinct, plumose. Upper floret ♀ stipitate, ± 2.5 mm long, lanceolate, obtuse, glabrous upwards; rachilla ± 1.4 mm long; callus densely bearded; lemma ovate-elliptic, obtuse, margins narrowly incurved, membranous, nerveless (-1-nerved), not keeled, glabrous or sparsely hairy in lower half; palea similar but widely incurved margins.

Type: INDIA: Tamil Nadu, Nilgiri Dist., Parthimund, 2300 m, July 11, 1970, J. L. Ellis 34627 (Holotype, CAL; Isotype, MH).

This variety is allied to var. *perpusilla* but differs in having broader culms, hirsute sheaths, broader blades, larger spikelets and panicle branches and nerveless and unkeeled upper lemma and palea; whereas var. *perpusilla* has capillary culms, glabrous sheaths, mostly filiform blades, smaller spikelets and panicle branches, and 2-nerved, 2-keeled and hairy upper lemma and palea. In var. *perpusilla*, the lower glume is 3-7-nerved but in this new variety lower glume is 1 or 3 nerved and when 3-nerved, lateral nerves present only in lower half.

ACKNOWLEDGEMENT

We are thankful to Dr. N. C. Majumdar who had kindly helped in Latin diagnosis of the new taxon.


VED PRAKASH

Botany Division, Central Drug Research Institute, Lucknow

AND

S. K. JAIN

Botanical Survey of India, Howrah



Coelachne perpusilla var. *nilagirica* Ved Prakash et Jain

Figs. 1-12: 1. Habit. 2. Spikelet. 3. Lower glume. 4. Upper glume. 5. Florets. 6. Lower lemma. 7. Lower palea. 8. Stamen. 9. Pistil. 10. Upper lemma. 11. Upper palea. (3, 4, 6, 7, 10, 11 all in ventral view). 12. Caryopsis. (J. L. Ellis 34627)